

Relativistic MHD Winds from Rotating Neutron Stars

N. Bucciantini¹, T.A. Thompson², J. Arons¹, E. Quataert¹

¹ Astronomy Department, University of California at Berkeley, 601 Campbell Hall, Berkeley, CA, 94720-3411, USA

² Department of Astrophysical Sciences, Peyton Hall, Ivy Lane, Princeton, NJ, 08544, USA

Abstract. We solve the time-dependent dynamics of axisymmetric, general relativistic MHD winds from rotating neutron stars. The mass loss rate is obtained self consistently as a solution of the MHD equations, subject to a finite thermal pressure at the stellar surface. Conditions are chosen to be representative of the neutrino driven phase in newly born magnetars, which have been considered as a possible engine for GRBs. We compute the angular momentum and energy losses as a function of σ and compare them with the analytic expectation from the classical theory of pulsar winds. We observe the convergence to the force-free limit in the energy loss and we study the evolution of the closed zone for increasing magnetization. Results also show that the dipolar magnetic field and the presence of a closed zone do not modify significantly the acceleration and collimation properties of the wind.

1. Introduction

Magnetically dominated outflows from stars and accretion discs, are a key element in understanding the evolution and properties of these objects. Magnetic outflows are ubiquitous in powering a variety of astrophysical systems, by converting the rotational energy into wind kinetic energy. Rotationally powered pulsars are among the best example of relativistic outflows produced by a fast compact rotator. The high magnetic field at the surface ($B \sim 10^{10-13}$ G) and short rotation period ($P \sim 0.001 - 1$ s) cause particles injection. If the charge density in the magnetosphere of the neutron star, due to pair creation, exceeds the Goldreich-Julian value then conditions can be modeled in terms of ideal RMHD.

Observations of the interaction of the pulsar wind with the surrounding SNR, tell us that far away from the source the wind is ultrarelativistic ($\gamma \sim 10^{4-6}$) and weakly magnetized (Arons 2004), in contrast with conditions inside the Light Cylinder, where $\gamma \sim 100$, and magnetization is high. This implies that the high Lorentz factor is achieved by almost completely converting magnetic energy into kinetic energy.

In this sense, pulsars constitute the prototype of relativistic accelerators: a plasma is assumed at the base with small outflow velocity (compared to the asymptotic velocity), embedded in a strong magnetic field. Magnetorotation then accelerates the plasma and converts the magnetic energy into relativistic motion.

In the case of pulsars, the plasma density at the surface of the neutron star, drops to very low values, and the magnetosphere is strongly magnetically dominated. Such conditions are close to force-free, and the equations that describe the flow dynamics become stiff, thus making impossible for present day codes to model such environment, in terms of MHD. Due to the presence of gaps at the foot points of open field lines, plasma is injected in the magnetosphere at a supersonic speed, quite differently from what happen in normal stars, where injection is subsonic.

However low magnetized neutron star winds are of interest primarily in understanding the physics of very young neutron stars. In the seconds after collapse and explosion, the neutron star is hot (surface thermal speed $\sim 0.1 - 0.03c$), and has an “extended atmosphere”. This proto-neutron star cools and contracts on its Kelvin-Helmholtz timescale ($\tau_{KH} \sim 10 - 100$ s), radiating its gravitational binding energy ($\sim 10^{53}$ ergs) in neutrinos of all species (Pons et al. 1999). The cooling epoch is accompanied by a thermal wind, driven by neutrino energy deposition, which emerges into the post-supernova-shock ejecta (Thompson et al. 2001).

After birth, the thermal pressure at the edge of the proto-neutron star surface decreases sharply as the neutrino luminosity decreases on a timescale τ_{KH} , as the star cools and deleptonizes. At some point during the cooling epoch the magnetic energy density will exceed the thermal pressure, and the wind region will become increasingly magnetically dominated. For magnetar-strength surface fields ($B_0 \sim 10^{14} - 10^{15}$ G) the magnetic field dominates the wind dynamics from just a few seconds after the supernova (Thompson 2003).

If magnetars are born with millisecond rotation periods (Thompson & Duncan 1993), their rotational energy can be very large ($\sim 2 \times 10^{52}$ ergs) relative to the energy of the supernova explosion ($\sim 10^{51}$ ergs) and be-




Fig. 1. Wind structure for a monopolar magnetic field, $\sigma = 200$, $\Omega = 0.14$. Left: poloidal magnetic field lines, and ratio $|B_\phi/B_r|$; Alfvénic (dashed) and fast (dotted) surfaces. Center: Lorentz factor, $\sim \sin(\theta)$. Right: energy flux in normalized units, $\sim \sin^2(\theta)$. From Bucciantini et al.(2006).

cause the spindown timescale can be short for larger value of the Alfvénic radius, these proto-magnetar winds have been considered as a mechanism for producing hyper-energetic supernovae (Thompson et al. 2004). Because the wind becomes increasingly magnetically-dominated, the outflow must eventually become relativistic. For this reason, proto-magnetar winds have also been considered as a possible central engine for long-duration GRBs (e.g., Thompson et al. 2004).

In the simple 1D radial model for relativistic winds from compact rotators by Michel (1969), solutions can be parameterized in terms of $\sigma = \Omega^2 \Phi^2 / \dot{M}$, where Ω is the rotation rate, Φ the magnetic flux and $\dot{M}$ the mass flux. σ is the maximum Lorentz factor the wind can achieve if all magnetic energy is converted into kinetic energy, and at the base of a pulsar it is $\sim 10^6$. The results by Michel (1969) show that the asymptotic Lorentz factor tends to $\sigma^{1/3}$. This is known as the $\sigma - \gamma$ problem (Arons 2004). If the flow is allowed to diverge more than radially then acceleration is more efficient (Begelman & Li 1994). The presence of a magnetic field can cause collimation of the flow along the rotation axis, and divergence along the equator, so in principle one might expect magnetic collimation to be also responsible for the acceleration of the wind. However when the flow speed reaches values close to c the Coulomb term in the electromagnetic force cannot be neglected and it balances the magnetic hoop stress. Solutions of the relativistic Grad-Shafranov equation for a monopolar field in the trans-fast regime (Beskin et al. 1998) have shown that acceleration is too slow and the asymptotic results of the Michel model apply also in 2D axisymmetric geometry, meaning that the flow remains radial.

However, no analytic multidimensional solution of the full relativistic MHD equations exists that extends from the surface of the neutron star to large distances. Ambiguities remain about injection conditions, which in the case of a newly-born hot magnetar, depend non-trivially on the temperature, gravity, and rotation. Even simplified

models consider only supersonic injection, but none focus on the possibility of a thermal wind at the base. Moreover the use of a monopolar field might prove not accurate, given that a more realistic dipole should be assumed in the close magnetosphere.

2. Wind acceleration

The first axisymmetric numerical simulations of the acceleration of a relativistic wind were presented by Bogovalov (1999), where the case of a zero pressure plasma in a monopolar magnetic field, with uniform injection conditions at the base of the star was investigated. Results show that as soon as $\gamma > 5$ collimation and acceleration of the wind become marginal.

Our recent investigation (Bucciantini et al. 2006) has focused on the study of the acceleration of an axisymmetric relativistic wind in the case the pressure is small compared with the rest mass energy density, but not negligible. If acceleration is efficient, the wind can reach high Lorentz factors ($\gamma \sim 100 - 300$) and kinetic luminosity ($\sim 10^{50}$ erg s $^{-1}$), perhaps enough to trigger a GRB. The relativistic MHD equations, together with Maxwell equations, are solved using the shock-capturing code for relativistic MHD developed by Del Zanna et al. (2003). The code has been modified to solve the equations in the Schwarzschild metric, in order to properly account the effects of gravity. In order to model the heating and cooling processes using an ideal gas equation of state we adopt an adiabatic coefficient $\Gamma = 1.1$ (almost isothermal wind). A hot atmosphere is assumed at the base of the star with $p/\rho c^2 \sim 0.01$. Density, pressure, rotation rate and the radial component of the magnetic field are fixed at the base of the neutron star, which is assumed to have a canonical radius $R_{\text{NS}} = 10$ km. The system is let to relax to a steady state configuration. We verify that it takes less than 5 rotations for the mass flux and energy losses to converge.




Fig. 2. Wind structure for a dipolar magnetic field, $\sigma = 20$, $\Omega = 0.14$. Left: poloidal magnetic field lines, and ratio $|B_\phi/B_r|$; Alfvénic (dashed) and slow (dotted) surfaces. Center: same as left, note the monopolar structure at larger radii. Fast (dotted) surface. Right: Lorentz factor, $\sim \sin(\theta)$. From Bucciantini et al.(2006).

In analogy with the 1D model, solutions in the 2D case can be parametrized by using the average value of σ over all stream lines. Results in the case of a monopolar magnetic field show that for $\sigma > 10$ the flow is close to radial, the terminal γ is $< \sigma$, pointing to inefficient acceleration. Moreover the energy and angular momentum fluxes rapidly approach the force-free values. Even if $v \ll c$ inside the Light Cylinder, asymptotic results agree with the flow structure found by Bogovalov (1999). Energy flux scales as $\sin^2(\theta)$, and $\gamma \propto \sin(\theta)$, in agreement with the solution of the exact monopole. Mass flux is also shaped by rotation. On the equator it appears to be 5 times higher than in the non rotating case. On the contrary due to collimation in the polar region the mass loss rate along the axis is about twice smaller than in the non rotating case. Such variation is due to the change in centrifugal support at different latitudes, and decreases as the period increases. Results suggest that a rapidly rotating magnetar will not be able to simultaneously produce a collimated and relativistic outflow. Relativistic velocities can be reached only for $\sigma \gg 1$, and in this case most of the energy would be concentrated in the equatorial plane instead that in a jet. Simulations have been extended to large distances ($\sim 10000 R_{NS}$) showing that for $\sigma > 10$ most of the energy is in the equatorial plane. We observe the formation of a collimated narrow jet, that is unresolved in our grid, but such structure appears not to be energetically significant.

We computed solutions also in the more realistic cases of dipolar magnetic field at the surface of the NS. Due to stability issues in numerical RMHD it was not possible to study cases with $\sigma > 20$. In the monopolar case, no current sheet is present. In the case of a dipole current sheets are present both at the edge of the closed zone and at the equator, and numerical resistivity is important. In particular in the equatorial plane, where the magnetic field vanishes, reconnection (due to numerical resistivity) takes place and plasmoids are formed. These plasmoids extend about 5-10 degrees around the equator, and cause




Fig. 3. Energy and angular momentum losses for a monopolar (upper panel) and dipolar (lower panel) magnetic field as a function of σ_0 (σ averaged on open field lines). Squares indicate cases with different rotation rate. Dashed-lines represent the low- σ analytic theory, dotted lines are the force-free limits. The solid line is a power law fit for the monopolar field. When normalized on open field lines, the results of the dipole follow the same curve of the monopole. From Bucciantini et al.(2006).

fluctuation of about 10-20% in the losses. Even if such values depend on numerical resistivity, this suggests that reconnection at the Y-point might manifest itself observationally.

To obtain steady solutions we suppressed resistivity at the equator, thus enhancing the instability of the code. Moreover, in the case of a dipole, the magnetic and flow surfaces are not aligned with the grid, and in this case inversion from conservative to primitive variables is not stable for $B^2/\rho c^2 > 100$.

Despite the presence of a closed zone, and the fact that within the Light Cylinder the field structure is close to a dipole, we found that solutions can again be parameterized in term of σ , now defined as an average done only on open streamlines. In this case we find that energy and angular momentum losses scale as a monopole with the same σ . Again as soon as $\sigma > 1$ solutions for the energy and angular momentum losses approach the force-free limit (Contopoulos et al. 1999), outside the Light Cylinder, and for $\sigma > 10$ the asymptotic structure resembles the exact monopole, with energy flux peaking at the equator and a higher equatorial Lorentz factor. Similar results are found also by Komissarov (2006). We are able to related the amount of open magnetic flux to the distance of the Y-point from the surface of the star. Extrapolating to the force-free limits, our result is in agreement with Gruzinov (2005).

Contrary to expectations (Contopoulos et al. 1999) we also find that the location of the Y-point is inside the Light Cylinder. This is due to the presence of a finite pressure plasma at the base of the star. In this case the extent of the closed zone is limited by the pressure equilibrium at the Y-point. We also verified that the ratio of the Y-point radius to the Light Cylinder R_Y/R_L drops with decreasing Ω , as one would expect for a braking index less than 3. This implies that even if for large σ the asymptotic wind structure is close to the force-free monopole solution, the spindown time can be shorter, and the total energy losses larger, that what is inferred by applying the dipole formula.

3. Conclusions

Recent numerical results have confirmed theoretical predictions about the acceleration of relativistic winds from neutron stars. The conversion of magnetic to kinetic energy appears to be minimal, the flow closely resembles the expectation from the exact monopole solution, there is no evidence for the formation of a collimated energetic jet. The use of a more realistic dipolar field does not seem to change significantly the wind properties far from the Light Cylinder. On the contrary the energy and angular momentum losses seem to depend only on the integrated value of σ , and, for $\sigma > 1$, on the amount of open magnetic flux.

Regarding the possibility that a magnetar could trigger a GRB, we conclude that, in ideal MHD, acceleration to relativistic speeds requires very high σ , much higher than what is thought to characterized newly-born magnetars. Moreover there is no evidence of energy collimation at high magnetization. Even if dissipation of the current sheet could be invoked to accelerate the wind, it is not obvious that a jet will form, it is more likely that the energy will still be concentrated in the equatorial plane, as is suggested by X-ray images of Pulsar Wind Nebulae.

Even if pulsar winds are not thermally driven at the base, they are cold, and the injection conditions depend on the physics of gaps, our results have several speculative implications for rotation powered pulsars, where σ is high. The fact that the Y-point is interior to the light cylinder (and the Alfvén radius), and that R_Y/R_L decreases as Ω decreases, suggests that observed braking indices less than 3 might be a consequence of pressure equilibrium. Even if globally, thermal pressure is not likely to be relevant for classical pulsars, magnetic dissipation at and near the Y point might cause pressure forces to be significant in this localized region.

The appearance of outwardly propagating plasmoids at and beyond the Y-point as a consequence of (numerical) magnetic dissipation raises the intriguing possibility that noise in pulsar spindown might arise from instability of the magnetospheric currents. Torque fluctuations might give rise to noise and limit cycles in the observed phase of pulsars' subpulses.

Acknowledgements. We gratefully acknowledge the support by the WE-Heraeus Foundation. This work has been supported by NSF grant AST-0507813, and by NASA grant NAG5-12031.

References

- Arons, J., 2004, 'Young Neutron Stars and Their Environments', I.A.U. Symposium, 218, 163
- Begelman, M. C., & Li, Z.-Y. 1994, ApJ, 426, 269
- Beskin, V. S., Kuznetsova, I. V., & Rafikov, R. R. 1998, MNRAS, 299, 341
- Bogovalov, S., 1999, A&A, 349, 1017
- Bucciantini, N., Thompson, T. A., Arons, J. et al., 2006, to be published in MNRAS, astro-ph/0602475
- Contopoulos, I., Kazanas, D., Fendt, C. 1999, ApJ, 511, 351
- Del Zanna, L., Bucciantini, N., Londrillo, P. 2003, A&A, 400, 397
- Gruzinov, A. 2005, Phys. Rev. Lett., 94
- Komissarov, S. S. 2006, MNRAS, 367, 19
- Michel, F. C. 1969, ApJ, 158, 727
- Pons, J. A., Reddy, S., Prakash, M., Lattimer, J. M., & Miralles, J. A. 1999, ApJ, 513, 780
- Thompson, C., & Duncan, R. C. 1993, ApJ, 408, 194
- Thompson, T. A., Burrows, A., Meyer, B. S. 2001, ApJ, 562, 887
- Thompson, T. A. 2003, ApJL, 585, L33
- Thompson, T. A., Chang, P., Quataert, E. 2004, ApJ, 611, 380